



HARVARD MEDICINE

NEWS ALERT

Harvard Medical School [Office of Public Affairs](#)

Simple chemical procedure augments therapeutic potential of stem cells

FINDINGS:

Researchers have developed a simple method for making a certain class of adult stem cells more therapeutically effective. By attaching a molecule called SLeX to the surface of human cells extracted from bone marrow, researchers have altered how the cells travel through vessels. This might enable the cells to more effectively reach sites of injury and replace damaged tissue.

RELEVANCE:

The cultured cells include mesenchymal stem cells, which can form fat cells, cartilage, bone, tendon and ligaments, muscle cells, and even nerve cells. When injected into the bloodstream of patients, mesenchymal stem cells can home to the site of an injury and replace damaged tissue. But just a fraction of the mesenchymal stem cells currently reach their target in clinical trials. One problem is that the cells lose their ability to roll along the walls of blood vessels when they grow and divide in a dish. The new technique should prevent cultured cells from drifting passively through the bloodstream, potentially enabling more of them to migrate out of the vessels into the surrounding tissue.



Dr. Jeffrey Karp

BOSTON, Mass. (Oct. 31, 2008) — Adult stem cells resemble couch potatoes if they hang out and divide in a dish for too long. They get fat and lose key surface proteins, which interferes with their movement and reduces their therapeutic potential. Now, via a simple chemical procedure, researchers have found a way to get these cells off the couch and over to their therapeutic target.

To do this, they simply added a molecule called SLeX to the surface of the cells. The procedure took just 45 minutes and restored an important biological function.


“Delivery remains one of the biggest hurdles to stem cell therapy,” explains senior author

Contact Us

Harvard Medical School
Office of Public Affairs
25 Shattuck Street
Boston, MA 02115

Telephone 617-432-0442
public_affairs@hms.harvard.edu

News Archives

Click  to subscribe to our news alerts. [What is RSS?](#)

Jeffrey Karp, an instructor at the Harvard-MIT Division of Health Sciences and Technology. “The blood stream offers a natural delivery vehicle, but stem cells don’t move through blood vessels normally after being expanded in culture. Our procedure promises to overcome this obstacle.”

These findings will be published online in the journal *Bioconjugate Chemistry* on Oct. 31.

In order for cells injected into the blood stream to be therapeutically useful, they need to take initiative to reach target tissues. But instead, cultured stem cells go with the flow. They move through the body quickly, carried by the current, which means they seldom contact the sides of blood vessels. Thus, they have fewer opportunities to escape into the surrounding tissue by squeezing between cells of the vessel wall. Adult stem cells must escape before they can colonize surrounding tissue and rebuild damaged structures.

In February of 2008, HMS associate professor Robert Sackstein (at Brigham and Women’s Hospital) and colleagues showed they could correct this problem by adding a particular molecule to the surface of adult stem cells. This molecule—a cousin of SLeX—formed temporary connections with proteins on the blood vessel wall, serving as a kind of weak tape. But Sackstein’s method involved enzymes, which made the chemistry complicated. Karp’s team achieved the same result without enzymes.

Karp lab postdoc Debanjan Sarkar simply flooded a dish of cells with three molecules—biotin, streptavidin, and SLeX—one after the other. The biotin and streptavidin anchored SLeX to the cell surface. Sarkar tweaked the concentrations of each molecule to maximize the cell’s ability to roll along the interior of the blood vessel, rather than getting lost in the flow. He also confirmed that the altered cells were still viable.

“The method is very simple,” says Sarkar, who is first author on the paper. “Plus, biotin and streptavidin work with many molecules, so labs can use this universal anchor we discovered to tackle other problems. They’re not limited to sticking SLeX on cells.”


The team worked with human cells extracted from the bone marrow. The cultures included mesenchymal stem cells (MSCs), which can form fat cells, cartilage, bone, tendon and ligaments, muscle cells, and even nerve cells. When injected into the bloodstream of patients, MSCs can home to the site of an injury and replace damaged tissue. But just a fraction of cultured MSCs currently reach their target in clinical trials. Karp’s procedure might improve their homing abilities.

Karp cautions that his lab’s discovery must be validated in animals, before doctors can apply it in the clinic. He’s collaborating with another lab to test the homing ability of the SLeX-dotted cells in mice.

“We need to confirm that this rolling behavior translates into increased homing and tissue repair,” explains Karp. “We may need to tweak the cells further.”

“This is definitely an approach that should be tried,” adds Pamela Robey, chief of the Craniofacial and Skeletal Diseases Branch of the National Institute of Dental and Craniofacial

DIAGRAM



Jeffrey Karp and colleagues prepared [this diagram](#) to illustrate how they increased the therapeutic potential of adult stem cells.

Research. Robey is working to reconstruct three-dimensional tissues with MSCs. "Jeff hasn't tested the altered MSCs inside animals, and that's really the gold-standard, but his *in vitro* data looks promising."

This research is supported by Brigham and Women's Hospital. The authors report no conflicts of interest.

Written by Alyssa Kneller

CITATION:

Bioconjugate Chemistry, Online Oct. 31, 2008

"Cell surface conjugation of Sialyl Lewis X induces a rolling response for human mesenchymal stem cells."

Debanjan Sarkar (1,2), Praveen Kumar Vemula (1,3,4), Grace Teo (1,2), Dawn Spelke (1,2), Rohit Karnik (5), Le Yee (1,2), Jeffrey Karp (1,2)

- (1) Harvard-MIT Division of Health Sciences and Technology
- (2) Harvard University
- (3) Department of Medicine, Brigham and Women's Hospital
- (4) Harvard Medical School
- (5) Massachusetts Institute of Technology, Mechanical Engineering

CONTACT:

Alyssa Kneller

public_affairs@hms.harvard.edu

617.432.0442

Harvard Medical School has more than 7,500 full-time faculty working in 11 academic departments located at the School's Boston campus or in one of 47 hospital-based clinical departments at 18 Harvard-affiliated teaching hospitals and research institutes. Those affiliates include Beth Israel Deaconess Medical Center, Brigham and Women's Hospital, Cambridge Health Alliance, Children's Hospital Boston, Dana-Farber Cancer Institute, Forsyth Institute, Harvard Pilgrim Health Care, Hebrew SeniorLife, Joslin Diabetes Center, Judge Baker Children's Center, Immune Disease Institute, Massachusetts Eye and Ear Infirmary, Massachusetts General Hospital, McLean Hospital, Mount Auburn Hospital, Schepens Eye Research Institute, Spaulding Rehabilitation Hospital, and VA Boston Healthcare System.

©2008 The President and Fellows of Harvard College